Please check the examination detail	ils below before ente	ering your candidate	information
Candidate surname		Other names	Thritishs the last
Centre Number Candida	te Number		Se Aijiele Altroom de 30a
Pearson Edexcel Int	ernation	al Advan	ced Level
Time 1 hour 30 minutes	Paper reference	WME	01/01
Mathematics			0 0
International Advanced Mechanics M1	d Subsidiar	y/Advanced	d Level
You must have: Mathematical Formulae and Stati	istical Tables (Ye	llow), calculato	Total Marks

Candidates may use any calculator permitted by Pearson regulations.

Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \,\mathrm{m\,s^{-2}}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶







1.




Figure 1

A particle P of weight 5 N is attached to one end of a light inextensible string. The other end of the string is attached to a fixed point O. The particle P is held in equilibrium by a force of magnitude F newtons. The direction of this force is perpendicular to the string and OP makes an angle of 60° with the vertical, as shown in Figure 1.

Find

(a) the value of F

(3)

Thins: Britishs the Barbon by John Med and

(b) the tension in the string.

(3)

Question 1 continued	hitos: Aritists to the Arto Con by	blank
	Troopport	od Web. an
		20)
		Q1
	(Total 6 marks)	



Leave blank

	Market Committee Com	blank
2.	A particle P has mass km and a particle Q has mass m . The particles are moving towards	
	each other in opposite directions along the same straight line when they collide directly	
	Immediately before the collision R has smood 24 and O has smood 4	
	Immediately before the conision, P has speed Su and Q has speed u .	, 100
	A particle P has mass km and a particle Q has mass m . The particles are moving towards each other in opposite directions along the same straight line when they collide directly, Immediately before the collision, P has speed $3u$ and Q has speed u . As a result of the collision, the direction of motion of each particle is reversed and the speed of each particle is halved.	* Ach app
	(a) Find the value of k .	
	(4)	
	(b) Find, in terms of m and u , the magnitude of the impulse exerted on Q in the collision. (3)	



Question 2 continued Villos Tarios T	blank
Question 2 continued This in the state of t	b _A 30a hebian

(Total 7 marks)	Q2



3. $\begin{array}{c|c}
 & 5 \text{ m} \\
\hline
 & D & C & B \\
\hline
 & 0.5 \text{ m}
\end{array}$

Figure 2

A beam ADCB has length 5 m. The beam lies on a horizontal step with the end A on the step and the end B projecting over the edge of the step. The edge of the step is at the point D where DB = 1.3 m, as shown in Figure 2.

When a small boy of mass 30 kg stands on the beam at C, where CB = 0.5 m, the beam is on the point of tilting.

The boy is modelled as a particle and the beam is modelled as a uniform rod.

(a) Find the mass of the beam.

(3)

1.3 m

A block of mass X kg is now placed on the beam at A.

The block is modelled as a particle.

(b) Find the smallest value of *X* that will enable the boy to stand on the beam at *B* without the beam tilting.

(3)

(c) State how you have used the modelling assumption that the block is a particle in your calculations.

(1)

Question 3 continued Thirds,	The state of the s
- "Mandantoon	The state of the s
	TO DE NOTE OF STREET OF ST
	Od Med app
	· Med alpo

	1



Overtion 2 continued	hips://dritists.tide.titooh.bd/	Leave
Question 3 continued	ttis _{lstuden} ,	
		6.
		a. Web. app

	A _{lbs://R}	blank
Question 3 continued	Ortos://Britishstatethroom/bg	
	······································	
		702.4°
		(S. ADD)
		Q3
	(Total 7 marks)	
	(Lotal / marks)	



At time $t = 0$, a small ball is projected vertically upwards from a point A who 24.5 m above the ground. The ball first comes to instantaneous rest at the point where $AB = 19.6$ m and first hits the ground at time $t = T$ seconds. The ball is modelled as a particle moving freely under gravity. (a) Find the value of T .	çh is
where $AB = 19.6$ m and first hits the ground at time $t = T$ seconds.	'VOQ
The ball is modelled as a particle moving freely under gravity.	
(a) Find the value of T.	
	(6)
(b) Sketch a speed-time graph for the motion of the ball from $t = 0$ to $t = T$ seconds.	
(No further calculations are needed in order to draw this sketch.)	(2)
	(2)



Question 4 continued	Alder Aritistestide Altoom by	blank
		, Da. Meb. app.



Mide	
h _{ttps://hritishs.htde.htro.ht/}	م
	X30'8'

	^{38:} /AritisIs III de III de Aritis II de II	Leave blank
Question 4 continued	Dritishsto.	
	*dentroom	
		Q. 4.
		ep. app
		0.4
		Q4
(Total	8 marks)	



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Hips://britistsindentoon/by/Qu/Nets/app.

Figure 3

A particle of mass m rests in equilibrium on a fixed rough plane under the action of a force of magnitude X. The force acts up a line of greatest slope of the plane, as shown in Figure 3.

The plane is inclined at an angle α to the horizontal, where $\tan \alpha = \frac{3}{4}$

The coefficient of friction between the particle and the plane is μ .

- When X = 2P, the particle is on the point of sliding up the plane.
- When X = P, the particle is on the point of sliding down the plane.

Find the value of μ .

l	o	١.



Question 5 continued	Alds: Aritists the control of the co	Leave blank
		A JOA WEB . A
		*\(\partial_{\partial\partial_{\partial_{\partial_{\partial_{\partial_{\partial_{\part



	· · · · · · · · · · · · · · · · · · ·	Leav
Question 5 continued	h _{IDS://Britishsthdentoombd3}	blanl
Question 5 continued	Ag _{lshsh} ,	

	ONA	
		8.4.
		(%)

Question 5 continued	hitos://thitististatetotooth/ba	blank
	*dentroon by	,
		Da. Web. app.
		Q5
	(Total 8 marks)	



6. [In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors.]

A particle P of mass $2 \log$ moves under the action of two forces, $(p\mathbf{i} + q\mathbf{j})N$ and $(2q\mathbf{i} + p\mathbf{j})N$, where p and q are constants.

Given that the acceleration of *P* is $(\mathbf{i} - \mathbf{j})$ m s⁻²

(a) find the value of p and the value of q.

(5)

(b) Find the size of the angle between the direction of the acceleration and the vector **j**.

(2)

At time t = 0, the velocity of P is $(3\mathbf{i} - 4\mathbf{j}) \,\mathrm{m} \,\mathrm{s}^{-1}$

At t = T seconds, P is moving in the direction of the vector $(11\mathbf{i} - 13\mathbf{j})$.

(c) Find the value of T.

(5)

Question 6 continued	Aldes: Abritishs de	Leave blank
		750g.
		web. app



uestion 6 continued	Augs:/Britishstateentroom/by
	- Jiroon d

Question 6 continued	Angs: Aritists the the too the basis	blank
		Od Web
		Mp.
		Q6
	(Total 12 marks)	



7.




Figure 4

A particle P of mass 4m lies on the surface of a fixed rough inclined plane.

The plane is inclined to the horizontal at an angle α where $\tan \alpha = \frac{3}{4}$

The particle *P* is attached to one end of a light inextensible string.

The string passes over a small smooth pulley that is fixed at the top of the plane. The other end of the string is attached to a particle Q of mass m which lies on a smooth horizontal plane.

The string lies along the horizontal plane and in the vertical plane that contains the pulley and a line of greatest slope of the inclined plane.

The system is released from rest with the string taut, as shown in Figure 4, and P moves down the plane.

The coefficient of friction between P and the plane is $\frac{1}{4}$

For the motion before Q reaches the pulley

(a) write down an equation of motion for Q,

(b) find, in terms of m and g, the tension in the string,

(7)

(4)

(c) find the magnitude of the force exerted on the pulley by the string.

(d) State where in your working you have used the information that the string is light.

(1)

	h _{ths}	Leave blank
Question 7 continued	Antiststatella	
	³ hden,	
	To the state of th	Pan bar
		×30,40,
		(d. dp)



Question 7 continued The both	Question 7 continued	*tos./Ariishs.	ŀ
		· · · · · · · · · · · · · · · · · · ·	
			303.46

Question 7 continued	h _{tos:/Ariists to de Atroon bos de}	blank
	"Udentroom b.	
		R. H. Ch. app
		Q7
(Tot	tal 13 marks)	



8. [In this question i and j are horizontal unit vectors directed due east and due and due blank blank respectively and position vectors are given relative to a fixed origin.]

**Valocity*(3i - 10j)km h⁻¹

At time t = 0, A is at the point with position vector (13i + 5j)km.

(a) Find \mathbf{r} in terms of t.

(2)

Another ship B moves with constant velocity (15i + 14j)km h⁻¹

At time t = 0, B is at the point with position vector $(3\mathbf{i} - 5\mathbf{j})$ km.

(b) Show that, at time t hours,

$$\overrightarrow{AB} = \left[(12t - 10)\mathbf{i} + (24t - 10)\mathbf{j} \right] \text{km}$$

(4)

Given that the two ships do not change course,

(c) find the shortest distance between the two ships,

(6)

(d) find the bearing of ship B from ship A when the ships are closest.

(2)





	A _{lly}	Leave
Overtion 9 continued	**************************************	blank
Question 8 continued	hips://Ariisishidentoon-ba	
	· uden	
	- Atoom .	
		D
		8.4°CX
		0.80p



						2			
						1	7	١	
							S		
							2		
							1		
						N			
						2		2	
						>			
							2		
	/					/	١	/	
						>			
						2		2	
						1		7	
						2			
١		è	Z	2	2	1	7	١,	
	Я	۲	7	۳			N		
	a		0	2					
		7	,	ę	?		2		
	/	á			à.		1		
	4	б							
	3	٠	۵	ú	ø		>		
2		2	₹	7		2		2	
		7		7	2	1			
	У	è	7				>		
		۵	й	ø	≺	>			
	А		ü	ь	œi		2		
		ļ	'n	ď	6	1	1	1	
	0		۲	3					
	я	ь.	2	S		κ			
		7	٠	7	۲,	2		2	
		è	4	\sim		Ν		/	
	Ŋ	-	7	e					
					₹	Þ			
				S	á	Κ	7	١,	
	М			۰	7		١		
		2	c						
	Я			c	2				
				7	٩		1		
						N		N	
		۵							
				7	₹	\rangle			
ζ	Ä	=	5	2	5	2	S	≷	
5	Š	7	2		Š	8	8	2	
5	Š	7	S	2	Ĭ	8	8	3	
>	Š		>		Š	8	5	2	
5	3		>		į	3	5	3	
5	3		2		į	3	3	3	
>	3		3			}	3	3	
	9	į	9		į	3	3	3	
			2		į		3	3	
			2		į			3	
			2		į		\ \ \	3	
			2					3	
								3	
								3	
								3	

			14 _{10/8.}	Leave blank
QX (Total 14 marks)	Question 8 continued		*Titishshider	
QX (Total 14 marks)			JIroo _{DD}	
QX (Total 14 marks)				Q. A.C. B.
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				
(Total 14 marks)				00
				Q8
TOTAL FOR PAPER: 75 MARKS				
END	E	TOTAL FOR PAPE ND	R: 75 MARKS	