Please check the examination details below before entering your candidate information					
Candidate surname	Other names				
Pearson Edexcel International Advanced Level	e Number Candidate Number				
Wednesday 15 M	Wednesday 15 May 2019				
Afternoon (Time: 1 hour 15 minutes) Paper Reference WCH06/01					
Chemistry Advanced Unit 6: Chemistry Laboratory Skills II					
Candidates must have: Scientific cald	Total Marks				

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶



P56132A
©2019 Pearson Education Ltd.
2/2/1/1/1//



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Answer ALL the questions. Write your answers in the spaces provided.	
1 This question is about five inorganic compounds (A, B, C, D and E).	
(a) Compounds A and B contain s-block elements.	
(i) In flame tests, A gave a yellow colour and B gave a yellow-red colour.	
Identify the s-block metal ions in ${\bf A}$ and ${\bf B}$.	(1)
	(1)
The metal ion in A	
The metal ion in B	
(ii) Give the colour of the precipitate formed when concentrated sodium hydroxide solution is added to an aqueous solution of B .	(1)
(iii) Compound A is a carbonate.	
Write an ionic equation for the reaction that takes place when dilute hydrochloric acid is added to an aqueous solution of A . Include state symbols.	(2)
(iv) Compound B is a halide.	
Identify, by name or formula, a reagent that may be used to test for halide ions in an aqueous solution of B .	(1)

2



DO NOT WRITE IN THIS AREA

(b) Compounds C , D and E are nitrates of d-block elements.	
Aqueous sodium hydroxide is added, drop by drop, until in excess to separate solutions of C , D and E .	
(i) Compound C forms an off-white precipitate which darkens on standing in ai	r.
Identify the precipitate, by name or formula, and explain why it darkens.	(2)
(ii) Compound D forms a white precipitate which dissolves in excess sodium hyo to form a colourless solution containing a complex ion.	droxide
Write the formula of this complex ion.	(1)
(iii) Compound E forms a green precipitate which does not dissolve in excess sodium hydroxide.	
The precipitate turns brown on standing in air.	
Identify, by name or formula, the compound E and the brown solid.	(2)
Compound E	
Brown solid	
(Total for Question 1 = 10 n	narks)
(Total for Question 1 = 10 ii	iiai K3)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2 This question is about three organic compounds: X, Y and Z. X, Y and Z have the same molecular formula, $C_6H_{12}O$.

The table shows the observations made in some chemical tests on X, Y and Z.

	Observations with reagent				
Compound	Sodium metal	Bromine water	sodium		lodine in aqueous sodium hydroxide
X	No change	No change	No change	Orange precipitate	Pale yellow precipitate
Υ	Effervescence	Decolourises	Turns green	No change	No change
Z	Effervescence	No change	No change	No change	No change

(1)	State what can be deduced about X from the positive test results.		
		(2))

(ii)	Name the functiona	I groups present in Y.
(11)	Name the functiona	i gioups piesent in i.

(2)



(a)

THIS AREA

(iii) Complete the equation for the reaction between **Z** and sodium metal. State symbols are not required.

(1)

$$C_6H_{12}O + Na \rightarrow$$

- (b) The **high** resolution proton nmr spectrum of compound **X** has only two peaks which are singlets with relative peak areas of 1:3.
 - (i) State what can be deduced from the presence of only two peaks in the nmr spectrum.

(ii) State what can be deduced from the fact that these peaks are singlets.

(1)

(1)

(iii) Use the nmr information, your answer to (a)(i) and the molecular formula to deduce the structure of **X**.

(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) Compound Y is straight-chained and does n	not have geometric or optical isomers.
---	---

(i) State what can be deduced from the fact that ${\bf Y}$ does not exist as geometric isomers.

(1)

(ii) State what can be deduced from the fact that **Y** does not have optical isomers.

(1)

(iii) Use information about **Y**, your answer to (a)(ii) and the molecular formula to deduce the structure of **Y**.

(1)

(d) (i) Deduce the type and classification of the functional group present in **Z**, using observations from the table.

(1)

(ii) Compound **Z** contains a five-membered carbon ring.

Deduce the structure of **Z** using this information, your deduction in (d)(i) and the molecular formula.

(1)

(Total for Question 2 = 13 marks)

6



THIS AREA

DO NOT WRITE IN

3 This question is about the analysis of iron supplements used to prevent or treat iron deficiency anaemia.

A student used the following procedure to analyse iron tablets containing iron in the form of hydrated iron(II) sulfate, FeSO₄.7H₂O.

Procedure

- Step **1** Grind up **two** iron tablets with a little dilute sulfuric acid using a pestle and mortar.
- Step 2 Transfer the resulting paste into a 100.0 cm³ volumetric flask. Rinse the apparatus used with dilute sulfuric acid, transferring all washings to the volumetric flask.
- Step **3** Add sufficient dilute sulfuric acid to the volumetric flask to make up the solution to exactly 100.0 cm³. Stopper the flask and invert it several times.
- Step **4** Using a pipette, transfer 10.0 cm³ of the solution to a conical flask and titrate it with 0.00500 mol dm⁻³ potassium manganate(VII) solution.
- Step **5** Repeat Step **4** until concordant results are obtained.

The overall equation for the reaction occurring in the titration is

$$MnO_4^-(aq) + 8H^+(aq) + 5Fe^{2+}(aq) \rightarrow Mn^{2+}(aq) + 5Fe^{3+}(aq) + 4H_2O(l)$$

(a) (i) Give the reason why the titration in Step **4** does **not** require the addition of an indicator.

(1)

(ii) Give the colour **change** at the end-point.

(1)



DO NOT WRITE IN THIS AREA

(b) The student decided to take the burette readings from the top of the liquid level rather than from the bottom of the meniscus.

Suggest the effect of this, if any, on the titre values. Justify your answer.

(2)

(c) Results of the titrations are given in the table.

Titration number	1	2	3	4
Burette reading (final) / cm ³	10.85	21.40	31.60	42.40
Burette reading (initial) / cm ³	0.00	10.85	21.40	32.10
Titre / cm ³	10.85			

(i) Complete the table and use the concordant values to calculate the mean titre.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(ii) Use your mean titre and information from the procedure to calculate the mass of hydrated iron(II) sulfate, FeSO₄.7H₂O, present in **one** iron tablet.

Give your answer to an appropriate number of significant figures.

(5)

- (d) The uncertainties in the burette and pipette measurements are $\pm 0.05\,\text{cm}^3$ and $\pm 0.06\,\text{cm}^3$ respectively.
 - Calculate which of these pieces of apparatus gives the greater percentage uncertainty in this experiment.

(2)

(Total for Question 3 = 13 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

4 This question is about the laboratory preparation of 3-methylbutyl ethanoate, an ester used as a banana flavouring in foods.

ethanoic acid 3-methylbutan-1-ol

3-methylbutyl ethanoate

Procedure

- Step **1** Add 7.5 cm³ of 3-methylbutan-1-ol, 10 cm³ of ethanoic acid and 2 cm³ of concentrated sulfuric acid to a round-bottom flask.
- Step 2 Add a few anti-bumping granules and heat the mixture under reflux for 35 minutes.
- Step **3** Transfer the cooled reaction mixture to a separating funnel. Add 30 cm³ of distilled water and washings from the flask. Shake the mixture, allow to separate and discard the aqueous layer.
- Step **4** Wash the organic layer with 15 cm³ of sodium hydrogencarbonate solution, releasing the build up of pressure. Discard the aqueous layer and repeat until the aqueous layer is slightly alkaline.
- Step **5** Transfer the organic layer to a boiling tube and dry with anhydrous magnesium sulfate.
- Step 6 Decant the organic layer into a clean round-bottom flask and distil.

 Collect the fraction boiling between 140 °C and 144 °C in a pre-weighed test tube.

Data

Compound	Molar mass / g mol ⁻¹	Density / g cm ⁻³	Boiling temperature / °C
3-methylbutan-1-ol	88.0	0.81	131
ethanoic acid	60.0	1.05	118
3-methylbutyl ethanoate	130.0	0.88	142



DO NOT WRITE IN THIS AREA

(a) State the purpose of the concentrated sulfuric acid and of the anti-bumping granules added to the round-bottom flask.	(2)
Concentrated sulfuric acid in Step 1	
Anti-bumping granules in Step 2	
(b) Draw a labelled diagram of the apparatus used to heat the reaction mixture under reflux in Step 2 .	(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) Draw a diagram of the separating funnel in Step 3, clearly labelling the aqueous and organic layers.

(2)

(d) Give the reason why the organic layer is washed with sodium hydrogencarbonate solution in Step **4** and suggest how the alkalinity of the aqueous layer should be confirmed.

(2

.....

12



DO NOT WRITE IN THIS AREA

e) Explain why the distillate is not collected below 140°C in Step 6 .	(2)
f) A student prepared 4.75 g of 3-methylbutyl ethanoate starting with 7.5 cm³ of 3-methylbutan-1-ol and excess ethanoic acid.	
(i) Calculate the percentage yield of 3-methylbutyl ethanoate.	(3)
	(3)
(ii) Give the main reason why the yield is significantly less than 100%.	
Do not consider errors in the experimental procedure or transfer losses.	(1)
(Total for Question 4 = 14 r	marks)
TOTAL FOR PAPER = 50 N	I A DI/C



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE

DO NOT WRITE IN THIS AREA DO NOT WRITE IN THIS AREA **BLANK PAGE** DO NOT WRITE IN THIS AREA 15

	0 (8) (18) 4.0	He helium 2	20.2	Ne	neon 10	39.9	Ar	argon 18	83.8	Ā	krypton 3 6	131.3	×e	xenon 54	[222]	R	radon 86		ted						
	_	(71)	19.0	ш	fluorine 9	35.5	ַ ז	chlorine 17	6.62	Ŗ	bromine 35	126.9	Н ;	iodine 53	[210]	Αt	astatine 85		Elements with atomic numbers 112-116 have been reported		175	Γn	lutetium 71	[257]	ځ
	9	(16)	16.0	0	oxygen 8	32.1	S	sulfur 16	79.0	Se	selenium 34	127.6	<u>e</u>	tellurium 52	[506]	8	polonium 84		116 have b	iticated	173	ΥÞ	ytterbium 70	[254]	<u>ڳ</u>
	22	(15)	14.0	z	nitrogen 7	31.0	₾ ;	phosphorus 15	74.9	As	arsenic 33	121.8	Sb	antimony 51	209.0	Bi	bismuth 83		mbers 112-	but not rully autnenticated	169	T	thulium 69	[526]	Þ₩
	4	(14)	12.0	U	carbon 6	28.1	Si	silicon 14	72.6	Ge	germanium 32	118.7	Sn	50 ti	207.2	Ъ	lead 82		atomic nui	but not 1	167	Ē	erbium 68	[253]	F
	m	(13)	10.8	В	boron 5	27.0	¥	aluminium 13	69.7	Ga	gallium 31	114.8	בַּ	indium 49	204.4	F	thallium 81		nents with		165	유	holmium 67	[254]	Es
ents								(12)	65.4	Zu	zinc 30	112.4	ਲ	cadmium 48	200.6	Нg	mercury 80		Elen		163	۵	dysprosium 66	[251]	Cf Es
Periodic Table of Elements								(11)	63.5	J C	copper 29	107.9	Ag	silver 47	197.0	PΠ	gold 79	[272]	Rg	roentgenium 111	159	Tb	terbium 65	[245]	쓢
le of								(10)	58.7	Ë	nickel 28	106.4	Pd	palladium 46	195.1	£	platinum 78	[271]	Ds	meitnerium damstadtium roentgenium	157	рg	gadolinium 64	[247]	E E
c Tab								(6)	58.9	ပ	cobalt 27	102.9	윤	rhodium 45	192.2	Ļ	iridium 77	[368]	¥	meitnerium 109	152	Eu	europium 63	[243]	Αm
riodi	1.0	H hydrogen 1						(8)	55.8	Fe	iron 26	101.1		ruthenium 44	190.2	Os	osmium 76	[277]	¥.	hassium 108	150	Sm	samarium 62	[242]	Pu
The Pe								(7)	54.9	Wn	manganese 25	[88]	ည	technetium 43	186.2	Re	rhenium 75	[264]		bohrium 107	[147]	Pm	promethium 61	[237]	Np Pu
广			mass	pol	umber			(9)	52.0	ъ	chromium manganese 24 25	95.9	Wo	molybdenum technetium 42 43	183.8		tungsten 74	[596]	Sg	seaborgium 106	144	PN	neodymium 60	238	⊃
		Key	relative atomic mass	atomic symbol	name atomic (proton) number			(2)	50.9	>	vanadium 23	92.9		niobium 41	180.9	٦a	tantalum 73	l		dubnium 105	141	P	prascodymium promethium samarium 59 60 61 62	[231]	Pa
			relati	ato	atomic			(4)	47.9	ï	titanium 22	91.2	Zr	zirconium 40	178.5	Ħ	hafnium 72	[261]	Æ.	rutherfordium 104	140	Çe	cerium 58	232	두
								(3)	45.0	Sc	scandium 21	88.9	>	yttrium 39	138.9	۲a*	lanthanum 57	[227]	A C*	actinium 89		Se			
	7	(2)	9.0	Be	beryllium 4	24.3	Mg	magnesium 12	40.1	Ca	calcium 20	87.6	S	strontium 38	137.3	Ba	barium 56	[526]	Ra	radium 88		* Lanthanide series	* Actinide series		
	-	ε	6.9	ב	lithium 3	23.0	Na	sodium 11	39.1	¥	potassium 19	85.5	₽	rubidium 37	132.9	C	caesium 55	[223]	<u>ት</u> (trancium 87		* Lanth	* Actini		