	alis below	before entering you	it pa/netidiadesituidentaoiem. wor
Candidate surname		Other r	names
Pearson Edexcel nternational dvanced Level	Centre	e Number	Candidate Number
Thursday 21	Ma	y 2020	
Afternoon (Time: 1 hour 30 minu	utes)	Paper Reference	te WME02/01
Mathematics International Advance Mechanics M2	ed Sub	osidiary/Ad	vanced Level

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \,\mathrm{m \, s^{-2}}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶



P65265A
©2020 Pearson Education Ltd.

	http://britishstudentroom.wordpre
1.	A particle of mass 2 kg is moving with velocity $(5\mathbf{i} + 3\mathbf{j})$ m s ⁻¹ when it receives an impulse INs, such that $\mathbf{I} = a\mathbf{i} + b\mathbf{j}$
	Immediately after receiving the impulse, the particle is moving with velocity $\lambda(\mathbf{i} + \mathbf{j}) \text{m s}^{-1}$, where λ is a constant.
	Given that the magnitude of I is $\sqrt{40}$, find the two possible impulses.
	(5)

	http://britishstudentroom.wordpress.leo bla
Question 1 continued	Ola
	Q1
	(Total 5 marks)



(9)

2. A truck of weight 9000 N is travelling up a hill on a straight road that is inclined at an angle θ to the horizontal, where $\sin \theta = \frac{1}{15}$

When the truck travels up the hill with the engine working at 3P watts, the truck is moving at a constant speed of $12 \,\mathrm{m\,s^{-1}}$

Later on, the truck travels down the hill along the same road, with the engine working at P watts. At the instant when the speed of the truck is $12 \,\mathrm{m\,s^{-1}}$, the acceleration of the truck . g

is $\frac{g}{20}$

The resistance to motion of the truck from non-gravitational forces is a constant force of magnitude R newtons in all circumstances.

Find (i) the value of P,

	···	.4	- 1	c D
((11)	the	value	of R .

	http://britishstudentroom.wordpre
estion 2 continued	
estion 2 continued	



	http://britishstudentroom.wordpress
uestion 2 continued	
destion 2 continued	

	http://britishstudentroom.wordpress
nestion 2 continued	
	Q
	(Total 9 marks)



3.




Figure 1

A uniform rod AB, of mass 25 kg and length 3 m, has end A resting on rough horizontal ground. The end B rests against a rough vertical wall.

The rod is in a vertical plane perpendicular to the wall.

The coefficient of friction between the rod and the ground is $\frac{4}{5}$

The coefficient of friction between the rod and the wall is $\frac{3}{5}$

The rod rests in limiting equilibrium.

The rod is at an angle of θ to the ground, as shown in Figure 1.

Find the exact value of $\tan \theta$.

(9)

DO NOT WRITE IN THIS AREA

	ntep.//orthonorusentroom.wordpre
	http://britishstudentroom.wordpre
estion 3 continued	



	http://britishstudentroom.wordpresslee
Question 3 continued	bia dia
4 10011011 0 101111111011	

	http://britishstudentroom.wordpress & bl
Question 3 continued	bl:
	Q3
	(Total 9 marks)



8*a* AВ aQ 2*a* R S 2a4a8*a* 4*a* UTDC

Figure 2

The uniform lamina L, shown shaded in Figure 2, is formed by removing the square PQRV, of side 2a, and the square RSTU, of side 4a, from a uniform square lamina ABCD, of side 8a. The lines QRU and VRS are straight. The side AD is parallel to PV and the side AB is parallel to PQ. The distance between AD and PV is a and the distance between AB and PQ is a. The centre of mass of L is at the point G.

(a) Show that the distance of G from the side AD is
$$\frac{42}{11}a$$
 (5)

The mass of L is M. A particle of mass kM is attached to L at C.

The lamina, with the attached particle, is freely suspended from B and hangs in equilibrium with BC making an angle of 45° with the horizontal.

(b)	Find the value of k .		
			(4)



	http://britishstudentroom.wordpre
estion 4 continued	
estion 4 continued	



	http://britishstudentroom.wordpres
estion 4 continued	
icition i continued	

http://britishstudentro		Ol.
	 	Q4



5. At time t seconds $(t \ge 0)$, a particle P has velocity $\mathbf{v} \, \mathbf{m} \, \mathbf{s}^{-1}$, where

$$\mathbf{v} = (3t^2 - 9t + 6)\mathbf{i} + (t^2 + t - 6)\mathbf{j}$$

(a) Find the acceleration of P when t = 3

(3)

When t = 0, P is at the fixed point O.

The particle comes to instantaneous rest at the point A.

(b) Find the distance *OA*.

(7)

	nttp://orthonstadentroom:wordpress.
uestion 5 continued	http://britishstudentroom.wordpress.l.
restion 5 continued	



	http://britishstudentroom.wordpress.ed
Question 5 continued	01
£44000000 0 000000000000000000000000000	

Question 5 continued		http://britishstudentroom.wordpressleen
	Question 5 continued	oran.
(Total 10 marks)		Q5
		(Total 10 marks)



DO NOT WRITE IN THIS AREA




Figure 3

A rough straight ramp is fixed to horizontal ground. The ramp has length 15 m and is inclined at an angle α to the ground, where $\tan \alpha = \frac{5}{12}$. The line AB is a line of greatest slope of the ramp, where A is at the bottom of the ramp, and B is at the top of the ramp, as shown in Figure 3.

A particle P of mass 6 kg is projected up the ramp with speed $14 \,\mathrm{m\,s^{-1}}$ from A in a straight line towards B. The coefficient of friction between P and the ramp is 0.25

(a) Find the work done against friction as P moves from A to B.

(3)

At the instant P reaches B, the speed of P is $v \text{ m s}^{-1}$. After leaving the ramp at B, the particle P moves freely under gravity until it hits the horizontal ground at the point C. Immediately before hitting the ground at C, the speed of P is $w \, \text{m s}^{-1}$

- (b) Use the work-energy principle to find
 - (i) the value of v,
 - (ii) the value of w.

(7)

	nttp://orthshistudentroom.wordpress.
uestion 6 continued	http://britishstudentroom.wordpress.k
restion o continueu	



s**leonn**∉ blank

	http://britishstudentroom.wor
Question 6 continued	

	http://britishstudentroom.wordpress & bl
Question 6 continued	bl
Question o continued	
	/T-4.140 1 \
	(Total 10 marks)



7. Particle A of mass 3m is moving in a straight line with speed 2u on a smooth horizontal surface. Particle A collides directly with particle B of mass m, which is moving along the same straight line and in the same direction as A.

Immediately before the collision, the speed of B is u.

As a result of the collision, the direction of motion of B is unchanged and the kinetic energy gained by B is $\frac{48}{25}mu^2$

(a) Find the coefficient of restitution between A and B.

(8)

After the collision, B hits a smooth fixed vertical wall that is perpendicular to the direction of motion of B. The coefficient of restitution between B and the wall is f.

Given that the speed of B immediately after first hitting the wall is equal to the speed of A immediately after its first collision with B,

(b) find the value of f.

(2)





estion 7 continued	http://britishstudentroom.wordpr
Ston / Continued	



s**leonn**∉ blank

	http://britishstudentroom.wor
Question 7 continued	

Question 7 continued	http://britisnstudentroom.wordpress	bla
		Ç
	(Total 10 marks)	



8. [In this question, the unit vectors \mathbf{i} and \mathbf{j} are in a vertical plane, with \mathbf{i} being horizontal and \mathbf{j} being vertically upwards.]




Figure 4

At time t = 0, a small ball is projected from a fixed point O on horizontal ground. The ball is projected from O with velocity $(p\mathbf{i} + q\mathbf{j}) \,\mathrm{m}\,\mathrm{s}^{-1}$, where p and q are positive constants. The ball moves freely under gravity.

At time t = 3 seconds, the ball passes through the point A with velocity $(8\mathbf{i} - 12\mathbf{j}) \,\mathrm{m} \,\mathrm{s}^{-1}$, as shown in Figure 4.

(a) Find the speed of the ball at the instant it is projected from O.

(5)

For an interval of T seconds the speed, $v \text{ m s}^{-1}$, of the ball is such that $v \leq 10$

(b) Find the value of *T*.

(4)

At the point B on the path of the ball, the direction of motion of the ball is perpendicular to the direction of motion of the ball at A.

(c) Find the vertical height of B above A.

(4)



	nttp://ortushstudentroom.wordpr
· 0 · · 1	http://britishstudentroom.wordpr
stion 8 continued	



	http://britishstudentroom.wordpress.k
Question 8 continued	b
guestion o continueu	

	http://britishstudentroom.wordpr
estion 8 continued	
estion o continued	



Question 8 continued	http://britishstudentroom.wordpr	ess!e@m blank
Zuestion o continueu		
		Q8
	(Total 13 marks)	
	TOTAL FOR PAPER: 75 MARKS	